

25. Seidman JD, Thomas RM. Multiple plexiform tumorlets of the uterus. *Arch Pathol Lab Med.* 1993;117(12):1255–1256.
26. Clement PB, Scully RE. Müllerian adenosarcoma of the uterus: a clinicopathologic analysis of 100 cases with a review of the literature. *Hum Pathol.* 1990;21(4):363–381.
27. Kanthan R, Senger JL. Uterine carcinosarcomas (malignant mixed müllerian tumours): a review with special emphasis on the controversies in management. *Obstet Gynecol Int.* 2011;2011:470795.
28. Fadare O. Perivascular epithelioid cell tumors (PEComas) and smooth muscle tumors of the uterus [letter]. *Am J Surg Pathol.* 2007;31(9):1454–1455.
29. Folpe AL, Mentzel T, Lehr HA, Fisher C, Balzer BL, Weiss SW. Perivascular epithelioid cell neoplasms of soft tissue and gynecologic origin: a clinicopathologic study of 26 cases and review of the literature. *Am J Surg Pathol.* 2005;29(12):1558–1575.
30. Vang R, Kempson RL. Perivascular epithelioid cell tumor ('PEComa') of the uterus: a subset of HMB-45-positive epithelioid mesenchymal neoplasms with an uncertain relationship to pure smooth muscle tumors. *Am J Surg Pathol.* 2002;26(1):1–13.
31. Dimmler A, Seitz G, Hohenberger W, Kirchner T, Faller G. Late pulmonary metastasis in uterine PEComa. *J Clin Pathol.* 2003;56(8):627–628.
32. Nogales FF, Isaac MA, Hardisson D, et al. Adenomatoid tumors of the uterus: an analysis of 60 cases. *Int J Gynecol Pathol.* 2002;21(1):34–40.
33. Al-Agha OM, Huwait HF, Chow C, et al. FOXL2 is a sensitive and specific marker for sex cord-stromal tumors of the ovary. *Am J Surg Pathol.* 2011;35(4):484–494.
34. Aron M, Gown AM, Balzer BL, et al. Comparative utility of novel nuclear markers steroidogenic factor (SF-1) and forkhead box L2 (FOXL2) in the diagnosis of sex cord stromal tumors (SCST) of the testis [abstract 791]. *Mod Pathol.* 2012;25(suppl 2):190A.