

REFERENCES:

- 1) Montgomery E, Fisher C: Paratesticular liposarcoma: a clinicopathologic study, Am J Surg Pathol 27:40–47, 2003
- 2) Hahn HP, Fletcher CD: Primary mediastinal liposarcoma: clinicopathologic analysis of 24 cases, Am J Surg Pathol 31:1868–1874, 2007.
- 3) Nascimento AF, McMenamin ME, Fletcher CD: Liposarcomas/atypical lipomatous tumors of the oral cavity: a clinicopathologic study of 23 cases, Ann Diagn Pathol 6:83–93, 2002
- 4) Cai YC, McMenamin ME, Rose G, et al: Primary liposarcoma of the orbit: a clinicopathologic study of seven cases, Ann Diagn Pathol 5:255–266, 2001.
- 5) Pontes FSC, de Souza LL, Vulcão ÉNC, de Abreu MC, Menezes SN, Rezende D, Pontes HAR. Liposarcoma of oral cavity: Systematic review of cases reported to date and analysis of prognostic factors. Head Neck. 2020 Sep;42(9):2626-2634. doi: 10.1002/hed.26160. Epub 2020 Apr 20. PMID: 32309895.
- 6) Pilotti S, Della Torre G, Mezzelani A, et al: The expression of MDM2/CDK4 gene product in the differential diagnosis of well differentiated liposarcoma and large deep-seated lipoma, Br J Cancer 82:1271–1275, 2000
- 7) Mariño-Enriquez A, Nascimento AF, Ligon AH, et al: Atypical spindle cell lipomatous tumor: clinicopathologic characterization of 232 cases demonstrating a morphologic spectrum, Am J Surg Pathol 41:234–244, 2017