

REFERENCES:

1. Matsuyama A, Shiba E, Umekita Y, Nosaka K, Kamio T, Yanai H, Miyasaka C, Watanabe R, Ito I, Tamaki T, Hayashi S, Hisaoka M. Clinicopathologic Diversity of Undifferentiated Sarcoma With BCOR-CCNB3 Fusion: Analysis of 11 Cases With a Reappraisal of the Utility of Immunohistochemistry for BCOR and CCNB3. *Am J SurgPathol*. 2017 Dec;41(12):1713-1721. doi: 10.1097/PAS.0000000000000934. PMID: 28877060.
2. Puls F, Niblett A, Marland G, Gaston CL, Douis H, Mangham DC, Sumathi VP, Kindblom LG. BCOR-CCNB3 (Ewing-like) sarcoma: a clinicopathologic analysis of 10 cases, in comparison with conventional Ewing sarcoma. *Am J SurgPathol*. 2014 Oct;38(10):1307-18. doi: 10.1097/PAS.0000000000000223. PMID: 24805859.
3. Machado I, Navarro L, Pellin A, et al. Defining Ewing and Ewinglike small round cell tumors (SRCT): the need for molecular techniques in their categorization and differential diagnosis. A study of 200 cases. *Ann DiagnPathol*. 2016;22:25–32.
4. Pierron G, Tirode F, Lucchesi C, Reynaud S, Ballet S, Cohen-Gogo S, Perrin V, Coindre JM, Delattre O. A new subtype of bone sarcoma defined by BCOR-CCNB3 gene fusion. *Nat Genet*. 2012 Mar 4;44(4):461-6. doi: 10.1038/ng.1107. PMID: 22387997.
5. Peters TL, Kumar V, Polikepahad S, Lin FY, Sarabia SF, Liang Y, Wang WL, Lazar AJ, Doddapaneni H, Chao H, Muzny DM, Wheeler DA, Okcu MF, Plon SE, Hicks MJ, López-Terrada D, Parsons DW, Roy A. BCOR-CCNB3 fusions are frequent in undifferentiated sarcomas of male children. *Mod Pathol*. 2015 Apr;28(4):575-86. doi: 10.1038/modpathol.2014.139. Epub 2014 Oct 31. PMID: 25360585; PMCID: PMC4385430.
6. Li WS, Liao IC, Wen MC, et al. BCOR-CCNB3-positive soft tissue sarcoma with round-cell and spindle-cell histology: a series of four cases highlighting the pitfall of mimicking poorly differentiated synovial sarcoma. *Histopathology*. 2016;69:792–801.
7. Brady EJ, Hameed M, Tap WD, Hwang S. Imaging features and clinical course of undifferentiated round cell sarcomas with CIC-DUX4